

Charged lepton electric dipole moments with the localized leptons and the new Higgs doublet in the two Higgs doublet model

E. O. Iltan *

Physics Department, Middle East Technical University
Ankara, Turkey

Abstract

We study the lepton electric dipole moments in the split fermion scenario, in the two Higgs doublet model, where the new Higgs scalars are localized around the origin in the extra dimension, with the help of the localizer field. We observe that the numerical value of the electron (muon, tau) electric dipole moment is at the order of the magnitude of 10^{-31} (10^{-24} , 10^{-22}) ($e - cm$) and this quantity is sensitive the new Higgs localization in the extra dimension.

*E-mail address: eiltan@newton.physics.metu.edu.tr

1 Introduction

The electric dipole moments (EDMs) of fermions are worthwhile to study since they are driven by the CP violating interactions. CP violation is carried by the complex Cabibo Kobayashi Maskawa (CKM) matrix elements in the quark sector and the possible lepton mixing matrix elements in the lepton sector, in the framework of the standard model (SM). The negligibly small theoretical values of the fermion EDMs in the SM forces one to search the physics beyond, such as multi Higgs doublet models (MHDM), supersymmetric model (SUSY), [1],..., etc..

There are various experimental results on the fermion EDMs in the literature and they read, $d_e = (1.8 \pm 1.2 \pm 1.0) \times 10^{-27} e\,cm$ [2], $d_\mu = (3.7 \pm 3.4) \times 10^{-19} e\,cm$ [3] and $|d_\tau| < (3.1) \times 10^{-16} e\,cm$ [4] respectively.

From the theoretical point of view, the extensive work has been done on the EDMs of fermions in various models, in the literature. The lepton electric dipole moments have been studied in the seesaw model, in [5]. The electron EDM has been predicted at the order of $10^{-32} e - cm$, in the 2HDM, including tree level flavor changing neutral currents (FCNC) and complex Yukawa couplings, in [6]. The work [7] is devoted to the fermion EDM moments in the SM with the inclusion of non-commutative geometry. In [8] ([9]), the electric dipole moments of fermions in the two Higgs doublet model with the inclusion of non-universal extra dimensions (in the split fermion scenario) have been analyzed. On the other hand, the quark EDMs have been estimated in several models [10] and the EDMs of nuclei, deuteron, neutron and some atoms have been predicted extensively [11].

In the present work, we study the lepton EDMs in the two Higgs doublet model in which the flavor changing (FC) neutral current vertices in the tree level are permitted and the CP violating interactions are carried by complex Yukawa couplings. In the calculations, we include the effects of a possible new dimension, in the split fermion scenario, where the hierarchy of fermion masses are coming from the overlap of the fermion Gaussian profiles in the extra dimension. In addition to the strong localization of the leptons, we consider that the new Higgs scalars are also localized around the the origin, in the extra dimension. These localizations are the result of the non vanishing couplings between leptons (new Higgs scalars) and the so called localizer field, which is odd under Z_2 reflection in the extra dimension. The split fermion scenario has been studied in several works in the literature [12]-[24]. In [12], the fermion mass hierarchy has been introduced by assuming that the fermions were located at different points in the extra dimensions, with exponentially small overlaps of their wave functions. The phenomenologically reliable locations of left and right handed components of fermions in the extra dimensions and their roles on the

mechanism of Yukawa hierarchies have been studied in [14]. [15] was devoted to prediction of the constraint on the split fermions in the extra dimensions by considering leptonic W decays and the lepton violating processes. The CP violation in the quark sector in the split fermion model was studied in [16] and in [17], the new configuration of split fermion positions in a single extra dimension and the physics of kaon, neutron and B/D mesons have been analyzed to find stringent bounds on the size of the compactification scale $1/R$. The shapes and overlaps of the fermion wave functions in the split fermion model have been studied in [18] and the rare processes in the split fermion scenario have been analyzed in [19]. [9] ([21],[22]) was related to the electric dipole moments of charged leptons (the radiative lepton flavor violating (LFV) decays, the LFV $Z \rightarrow l_i l_j$ decays) in the split fermion scenario. The work [23] was devoted to the branching ratios of the radiative LFV decays in the split fermion scenario, with the assumption that the new Higgs doublet is restricted to the 4D brane or to the thin bulk in one and two extra dimensions, in the framework of the two Higgs doublet model. Recently, the Higgs localization in the split fermion models has been studied in [24].

The present work is devoted to the EDMs of charged leptons in the 2HDM with the assumption that they have Gaussian profiles in the extra dimension, similar to the previous work [9]. However, we also consider that the new Higgs scalars are localized in the extra dimension with the help of the so called localizer field. The idea of the localization of the SM Higgs, using the localizer field, has been studied in [24]. Here, we assume that the new Higgs scalars are localized in the extra dimension and the SM Higgs has a constant profile. The localization of the new Higgs scalars depends strongly on the strength of the coupling of the localizer to the new Higgs scalar. Here we take this coupling small and study the sensitivity of the lepton EDMs on it. Furthermore, we analyze the compactification scale dependence of lepton EDMs. In the numerical calculations, we observe that the lepton EDMs are sensitive to the new Higgs localization in the extra dimension and we estimate that the numerical values of electron (muon, tau) EDM is at the order of the magnitude of 10^{-31} (10^{-24} , 10^{-22}) e-cm, even for the large values of the compactification scale $1/R$ and for the intermediate values of Yukawa couplings.

The paper is organized as follows: In Section 2, we present EDMs of charged leptons in the split fermion scenario, with the localization of new Higgs scalars in the extra dimension, in the 2HDM. Section 3 is devoted to discussion and our conclusions. In Appendix section we present the derivation of the KK modes of new Higgs fields and their masses in the case that they are coupled to the localizer field.

2 The effect of the localization of new Higgs scalars on the electric dipole moments of charged leptons, in the split fermion scenario, in the two Higgs doublet model.

The existence of the fermion EDM is the signal for the CP violation, since it is the result of the CP violating fermion-fermion-photon effective interaction. In the framework of the SM, the complex CKM matrix (possible lepton mixing matrix) is the possible source of this violation for quarks (for charged leptons). Since the theoretical estimated numerical values of the fermion EDMs are considerably small, there is a need to go beyond the SM to get additional contributions, in order to enhance their numerical values. The multi Higgs doublet models are among the candidates for this enhancement, and, in the present work, we take the 2HDM, which allows the FCNC at tree level and includes the complex Yukawa couplings as a source of CP violation. On the other hand, with the addition of extra dimensions, there exist new contributions sensitive to the compactification scale $1/R$ where R is the compactification radius of the extra dimension. Here, we take the effects of extra dimensions into account with the assumption that the hierarchy of lepton masses are coming from the lepton Gaussian profiles in the extra dimensions, so called the split fermion scenario. This localization is obtained by the nonzero coupling of the localizer scalar field ϕ_L to the lepton fields. Furthermore, we assume that this field couples also to the new Higgs doublet and localizes the new Higgs scalars around the origin $y = 0$, where y is the coordinate of the extra dimension (see Appendix).

The Yukawa Lagrangian which creates the lepton EDM in a single extra dimension, respecting the split fermion scenario, reads:

$$\mathcal{L}_Y = \xi_{5ij}^E \bar{\hat{l}}_{iL} \phi_2 \hat{E}_{jR} + h.c. , \quad (1)$$

where L and R denote chiral projections $L(R) = 1/2(1 \mp \gamma_5)$, ϕ_2 is the new scalar doublet. Here $\hat{l}_{iL}$ ($\hat{E}_{jR}$), with family indices i, j , are the zero mode ¹ lepton doublets (singlets) with Gaussian profiles in the extra dimension y and they read

$$\begin{aligned} \hat{l}_{iL} &= N e^{-(y-y_{iL})^2/2\sigma^2} l_{iL}, \\ \hat{E}_{jR} &= N e^{-(y-y_{jR})^2/2\sigma^2} E_{jR}, \end{aligned} \quad (2)$$

with the normalization factor $N = \frac{1}{\pi^{1/4}\sigma^{1/2}}$. l_{iL} (E_{jR}) are the lepton doublets (singlets) in four dimensions. Here the parameter σ is the Gaussian width of the leptons with the property

¹In our calculations, we take only zero mode lepton fields.

$\sigma \ll R$, and $y_{i(L,R)}$ are the fixed position of i^{th} left (right) handed lepton in the fifth dimension².

The Higgs doublets ϕ_1 and ϕ_2 are chosen as

$$\phi_1 = \frac{1}{\sqrt{2}} \left[\begin{pmatrix} 0 \\ v + H^0 \end{pmatrix} + \begin{pmatrix} \sqrt{2}\chi^+ \\ i\chi^0 \end{pmatrix} \right] ; \phi_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}H^+ \\ H_1 + iH_2 \end{pmatrix} . \quad (4)$$

so that their vacuum expectation values read:

$$\langle \phi_1 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix} ; \langle \phi_2 \rangle = 0 . \quad (5)$$

This leads to the possibility of collecting SM (new) particles in the first (second) doublet. In this case H_1 and H_2 are the mass eigenstates h^0 and A^0 respectively since no mixing occurs between two CP-even neutral bosons H^0 and h^0 at tree level.³

Now we take that the SM Higgs has a constant profile in the extra dimension and the new Higgs field ϕ_2 , which has the main role in the existence of the charged lepton EDM, couples to the localizer ϕ_L with a small coupling g (see the Appendix section for details). The coupling of the new Higgs doublet to the localizer results in the localization of the Higgs scalars, with the corresponding KK modes, around the origin, and brings modified Yukawa interactions in four dimensions. The neutral CP even and odd scalar fields S ($S = h^0, A^0$), are expanded into their KK modes as $S(x, y) = \sum_n h_n(y) S^{(n)}(x)$ (see the Appendix section) and to obtain the lepton-Higgs interaction coupling in four dimensions we need to integrate the combination $\bar{f}_{iL(R)} S^{(n)}(x) h_n(y) \hat{f}_{jR(L)}$, appearing in the part of the Lagrangian (eq. (1)), over the fifth dimension. Using the KK basis obtained (see eq. (26) and (28)), we get

$$\int_{-\pi R}^{\pi R} dy \bar{f}_{iL(R)} S^{(n)}(x) h_n(y) \hat{f}_{jR(L)} = V_{LR(RL)ij}^n \bar{f}_{iL(R)} S^{(n)}(x) f_{jR(L)} , \quad (7)$$

²The positions of left handed and right handed leptons are obtained by taking the observed masses into account [14]. The idea is that the lepton mass hierarchy is due to the relative positions of the Gaussian peaks of the wave functions located in the extra dimension [12, 14]. By assuming that the lepton mass matrix is diagonal, one possible set of locations for the lepton fields read (see [14] for details)

$$P_{li} = \sqrt{2}\sigma \begin{pmatrix} 11.075 \\ 1.0 \\ 0.0 \end{pmatrix} , \quad P_{ei} = \sqrt{2}\sigma \begin{pmatrix} 5.9475 \\ 4.9475 \\ -3.1498 \end{pmatrix} . \quad (3)$$

³Here we consider the Higgs potential term $V(\phi_1, \phi_2, \phi_L)$ as,

$$V(\phi_1, \phi_2, \phi_L) = c_1(\phi_1^\dagger \phi_1 - v^2/2)^2 + c_2(\phi_2^\dagger \phi_2)^2 + \frac{1}{2} g \phi_2^\dagger \phi_2 \phi_L^2 , \quad (6)$$

with constants c_i , $i = 1, 2$, where ϕ_L is the localizer singlet field (see Appendix section for details). In this case, the first Higgs doublet does not couple to the second one and to the localizer singlet field. With this assumption, the two Higgs doublets do not mix and the SM (new) particles are placed in the first (second) doublet. If the first and the second doublets couple and the second one acquires non zero vacuum expectation value, the mixing appears between two doublets and it needs further analysis.

where the factor $V_{LR(RL)ij}^n$ reads

$$V_{LR(RL)ij}^n = V_{LR(RL)ij}^0 c_n^{(i,j)}, \quad (8)$$

and the function $V_{LR(RL)ij}^0$ is

$$V_{LR(RL)ij}^0 = \frac{g^{1/8} e^{-(y_{iL(R)} - y_{jR(L)})^2 / 4\sigma^2}}{\sqrt{\sigma} \pi^{1/4} \sqrt{\text{Erf}[\pi g^{1/4} \frac{R}{\sigma}]}}. \quad (9)$$

Here, the fields f_{iL} , f_{jR} are the four dimensional lepton fields and the function $\text{Erf}[z]$ is the error function, which is defined as

$$\text{Erf}[z] = \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2} dt. \quad (10)$$

The functions $c_n^{(i,j)}$ in eq. (8) are calculated in the case that the coupling g is non zero and small:

$$\begin{aligned} c_n^{(i,j)} &= \frac{(-1)^n 2^{3n+\frac{1}{2}} \sqrt{\pi} (1 - \frac{g}{4})^n (2 + \sqrt{g})^{-\frac{4n+1}{2}}}{\sqrt{(2n)!} \Gamma[\frac{1}{2} - n]} e^{f_{LR(RL)ij}} \\ &\times {}_1F_1[-n, \frac{1}{2}, \frac{\sqrt{g} (y_{iL(R)} + y_{jR(L)})^2}{(4 - g) \sigma^2}], \end{aligned} \quad (11)$$

where

$$f_{LR(RL)ij} = -\frac{\sqrt{g} (y_{iL(R)} + y_{jR(L)})^2}{4 (2 + \sqrt{g}) \sigma^2}. \quad (12)$$

The function ${}_1F_1[a; b; z]$ appearing in eq. (11) is the hypergeometric function

$${}_1F_1[a; b; z] = \sum_{k=0}^{\infty} (a)_k (b)_k z^k / k!, \quad (13)$$

where $(d)_k = \frac{\Gamma[d+k]}{\Gamma[d]}$. Finally, we can define the Yukawa couplings in four dimensions as

$$\xi_{ij}^E ((\xi_{ij}^E)^\dagger) = V_{LR(RL)ij}^0 \xi_{5ij}^E ((\xi_{5ij}^E)^\dagger), \quad (14)$$

where ξ_{5ij}^E are the Yukawa couplings in five dimensions (see eq. (1))⁴.

The effective EDM interaction for a charged lepton f is given by

$$\mathcal{L}_{EDM} = id_f \bar{f} \gamma_5 \sigma^{\mu\nu} f F_{\mu\nu}, \quad (15)$$

⁴In the following we use the dimensionful coupling $\bar{\xi}_N^E$ in four dimensions, with the definition $\xi_{N,ij}^E = \sqrt{\frac{4G_F}{\sqrt{2}}} \bar{\xi}_{N,ij}^E$ where N denotes the word "neutral".

where $F_{\mu\nu}$ is the electromagnetic field tensor, ' d_f ' is EDM of the charged lepton and it is a real number by hermiticity. With the assumption that the CP violating EDM interaction comes from the complexity of the Yukawa couplings due to the new Higgs scalars⁵, the f -lepton EDM ' d_f ' ($f = e, \mu, \tau$) can be calculated as a sum of contributions coming from neutral Higgs bosons h_0 and A_0 (see Fig. 1),

$$d_f = -\frac{i G_F}{\sqrt{2}} \frac{e}{32\pi^2} \frac{Q_\tau}{m_\tau} \left((\bar{\xi}_{N,l\tau}^{E*})^2 - (\bar{\xi}_{N,\tau l}^E)^2 \right) \left(c_0(f, \tau) c'_0(f, \tau) (F_1(y_{h_0}) - F_1(y_{A_0})) \right. \\ \left. + \sum_{n=1}^{\infty} c_n(f, \tau) c'_n(f, \tau) (F_1(y_{h_0}^n) - F_1(y_{A_0}^n)) \right), \quad (16)$$

for $f = e, \mu$ and

$$d_\tau = -\frac{i G_F}{\sqrt{2}} \frac{e}{32\pi^2} \left\{ \frac{Q_\tau}{m_\tau} \left((\bar{\xi}_{N,\tau\tau}^{E*})^2 - (\bar{\xi}_{N,\tau\tau}^E)^2 \right) \left(c_0^2(\tau, \tau) (F_2(r_{h_0}) - F_2(r_{A_0})) \right. \right. \\ \left. \left. + \sum_{n=1}^{\infty} c_n^2(\tau, \tau) (F_2(r_{h_0}^n) - F_2(r_{A_0}^n)) \right) \right. \\ \left. - Q_\mu \frac{m_\mu}{m_\tau^2} \left((\bar{\xi}_{N,\mu\tau}^{E*})^2 - (\bar{\xi}_{N,\tau\mu}^E)^2 \right) \left(c_0(\mu, \tau) c'_0(\mu, \tau) (r_{h_0} \ln(z_{h_0}) - r_{A_0} \ln(z_{A_0})) \right. \right. \\ \left. \left. + \sum_{n=1}^{\infty} c_n(\mu, \tau) c'_n(\mu, \tau) (r_{h_0}^n \ln(z_{h_0}^n) - r_{A_0}^n \ln(z_{A_0}^n)) \right) \right\}, \quad (17)$$

for $f = e, \mu, \tau$. Here the functions $F_1(w)$, $F_2(w)$ read

$$F_1(w) = \frac{w(3 - 4w + w^2 + 2 \ln w)}{(-1 + w)^3}, \\ F_2(w) = w \ln w + \frac{2(-2 + w)w \ln \frac{1}{2}(\sqrt{w} - \sqrt{w-4})}{\sqrt{w(w-4)}}, \quad (18)$$

with $y_S = \frac{m_\tau^2}{m_S^2 + 2\beta}$, $y_S^n = \frac{m_\tau^2}{m_S^2 + 2(4n+1)\beta}$, $r_S = \frac{1}{y_S}$, $r_S^n = 1/y_S^n$, $z_S = \frac{m_\mu^2}{m_S^2 + 2\beta}$, $z_S^n = \frac{m_\mu^2}{m_S^2 + 2(4n+1)\beta}$, Q_τ , Q_μ are charges of τ and μ leptons respectively. Notice that the functions $c_n^{(i)}(i, j)$ ⁶ are defined in eq. (11). In eq. (16) we take into account only the internal τ -lepton contribution respecting our assumption that the Yukawa couplings $\bar{\xi}_{N,ij}^E$, $i, j = e, \mu$, are small compared to $\bar{\xi}_{N,\tau i}^E$, $i = e, \mu, \tau$ due to the possible proportionality of the Yukawa couplings to the masses of leptons in the vertices. In eq. (17), we present also the internal μ -lepton contribution, which

⁵We do not consider the possible effects due to the CKM type lepton mixing matrix and take only zero mode lepton fields.

⁶If the second Higgs doublet is localized around $y = y_0$, the zero mode solution reads $h_0(y) = N' e^{-\beta(y-y_0)^2}$ (see eq. (26)), and, if y_0 is away from the origin, $y_0 \rightarrow \pi R$, the coefficients $c_n^{(i)}(i, j)$ tends to zero. This is the case where the new Higgs contribution vanishes.

can be neglected numerically. Notice that, we make our calculations in arbitrary photon four momentum square q^2 and take $q^2 = 0$ at the end.

Finally, in our calculations, we choose the Yukawa couplings complex and we used the parametrization

$$\bar{\xi}_{N,\tau f}^E = |\bar{\xi}_{N,\tau f}^E| e^{i\theta_f}. \quad (19)$$

Therefore, the Yukawa factors in eqs. (16) and (17) can be written as

$$((\bar{\xi}_{N,f\tau}^{E*})^2 - (\bar{\xi}_{N,\tau f}^E)^2) = -2i \sin 2\theta_f |\bar{\xi}_{N,\tau f}^E|^2 \quad (20)$$

where $f = e, \mu, \tau$. Here θ_f is the CP violating parameter which is the source of the lepton EDM.

3 Discussion

The fermion EDM is the result of the existence of the CP violation and arises from the fermion-fermion-photon effective interaction. In the present work, we study the charged lepton EDMs in the 2HDM with FCNC at tree level and we assume that the CP phase is coming from the complex Yukawa couplings, driving the lepton-lepton-new Higgs scalar vertices. Furthermore, we take into account the effects of a single extra dimension in the so called split fermion scenario where the hierarchy of lepton masses are coming from the lepton Gaussian profiles in the extra dimension. In this scenario, the penetration of leptons into the bulk is ensured by the non-zero coupling of leptons with a background scalar field, namely the localizer field, ϕ_L , which has a vacuum expectation value, depending on the coordinate of the extra dimension and centered at the origin. In addition to this, we consider that the new Higgs doublet also couples to the localizer weakly and the corresponding Higgs scalars localize around the origin $y = 0$. On the other hand, the SM Higgs H^0 is taken as uniform in the extra dimension and its vacuum expectation value is responsible for the generation of the masses. Notice that the mass term, which is modulated by the mutual overlap of lepton wavefunctions, is obtained by integrating the operator $H^0 \bar{f} \hat{f}$ over extra dimensions, where f denotes lepton. This is the idea of fixing the positions of left (right) handed leptons in the extra dimensions (see [14] for details). Since the lepton-lepton-new scalar vertices creates the CP violating EDM interaction, besides zero modes, the KK modes of leptons and Higgs scalars have additional contributions. Here, we include the effects of Higgs KK modes, however, we ignored the lepton KK mode contributions to the EDMs of leptons, by expecting that, for the large values of the compactification scale, their effects on the physical parameters are suppressed.

In the numerical calculations, the free parameters coming the model used should be fixed by using the present experimental results. The first set of free parameters is the four dimensional leptonic complex couplings $\bar{\xi}_{N,ij}^E$, $i, j = e, \mu, \tau$. We consider the Yukawa couplings $\bar{\xi}_{N,ij}^E$, $i, j = e, \mu$, as smaller compared to $\bar{\xi}_{N,\tau i}^E$ $i = e, \mu, \tau$ and we assume that $\bar{\xi}_{N,ij}^E$ is symmetric with respect to the indices i and j . In the case that no extra dimension exists, the experimental uncertainty, 10^{-9} , in the measurement of the muon anomalous magnetic moment [25] can be used to estimate the upper limit of $\bar{\xi}_{N,\tau\mu}^E$, since the new physics effects can not exceed this uncertainty and this limit was predicted as 30 GeV (see [26] and references therein) . Using this upper limit and the experimental upper bound of BR of $\mu \rightarrow e\gamma$ decay, $\text{BR} \leq 1.2 \times 10^{-11}$, the coupling $\bar{\xi}_{N,\tau e}^E$ is restricted in the range, $10^{-3} - 10^{-2} \text{ GeV}$ (see [6]). In our calculations, we choose the numerical values of the couplings $\bar{\xi}_{N,\tau\mu}^E$ ($\bar{\xi}_{N,\tau e}^E$) not larger than 30 GeV (around 10^{-3} GeV). For the coupling $\bar{\xi}_{N,\tau\tau}^E$, we use the numerical values which are greater than $\bar{\xi}_{N,\tau\mu}^E$, since we have no explicit restriction region. For the CP violating parameter which drives the EDM interaction we choose the intermediate value, $\sin \theta_{e(\mu,\tau)} = 0.5$

The second set of free parameters is coming from the split fermion scenario where the hierarchy of lepton masses are due to the lepton Gaussian profiles in the extra dimension. Here, we have a free parameter $\rho = \sigma/R$, where σ is the Gaussian width of the fermions and R is the compactification radius. There are various predictions on the compactification radius in the literature. The lower bound for inverse of the compactification radius was estimated as $\sim 300 \text{ GeV}$ [27] in the case of universal extra dimension scenario. The direct limits from searching for KK gauge bosons imply $1/R > 800 \text{ GeV}$, the precision electro weak bounds on higher dimensional operators generated by KK exchange place a far more stringent limit $1/R > 3.0 \text{ TeV}$ [28] and, from $B \rightarrow \phi K_S$, the lower bounds for the scale $1/R$ have been obtained as $1/R > 1.0 \text{ TeV}$, from $B \rightarrow \psi K_S$ one got $1/R > 500 \text{ GeV}$, and from the upper limit of the BR, $\text{BR}(B_s \rightarrow \mu^+\mu^-) < 2.6 \times 10^{-6}$, the estimated limit was $1/R > 800 \text{ GeV}$ [19]. On the other hand, there exist different phenomenologically reliable set of locations of lepton fields in the extra dimension, even if they are predicted by using the lepton masses. Notice that we used the set ⁷ predicted by the work [14]. Finally, the coupling g , which exists since we consider that the new Higgs doublet also couples to the localizer, is another free parameter of the model used. Here, we assume that this coupling is in a region such that the new Higgs scalars are localized around the origin as much as possible. Notice that the zero mode Higgs boson h^0 and A^0 masses $m_{0,h^0(A^0)}^2 = m_{h^0(A^0)}^2 + 2\beta$ (with $\beta = \sqrt{g}\mu^2$) depend on the coupling g

⁷The numerical results of physical parameters depends on choice of the set of locations of the lepton fields.

and the additional term comes from the localization effect, which should not be large ⁸.

Now, we start to estimate the charged lepton EDMs and study the coupling g and the parameter ρ dependence of these physical quantities.

In Fig. 2, we plot the electron EDM d_e with respect to the coupling g for $1/R = 1000 \text{ GeV}$ and $\bar{\xi}_{N,\tau e}^E = 0.001 \text{ GeV}$. Here the lower-upper solid (dashed) line represents the EDM, due to the zero mode Higgs-KK mode Higgs included contribution for the parameter $\rho = 0.001$ ($\rho = 0.0005$). The sensitivity of EDM to the coupling g is strong. The addition of KK mode contributions enhance the EDM weakly, especially for the parameter $\rho = 0.001$. The electron EDM is at most at the order of the magnitude of $10^{-31} - 10^{-32} (e - cm)$, for $g \sim 10^{-14}$ and $\rho = 0.001$ ($\rho = 0.0005$). The enhancement in EDM for the small values of the coupling g (and also for the large values of the width σ) is due the fact that the masses of zero (and also KK modes) (see eq. (29)) are smaller compared to the ones for the large values of the coupling g (and also for the small values of the width σ). In the case that all the 2HDM particles are confined into the 4D brane and the extra dimension effects are switched off, the electron EDM d_e is at the order of $10^{-30} (e - cm)$, which is the numerical value to be reached with smaller coupling g and the weak localization of the new Higgs doublet, in the present scenario.

Fig. 3 is devoted to the compactification scale $1/R$ dependence of d_e , for $g = 10^{-13}$ and $\bar{\xi}_{N,\tau e}^E = 0.001 \text{ GeV}$. Here the lower-upper solid (dashed) line represents the d_e , due to the zero mode Higgs-KK mode Higgs included contribution for the parameter $\rho = 0.001$ ($\rho = 0.0005$). The sensitivity of d_e to $1/R$ is strong and it enhances with the decreasing values of $1/R$, as expected. The effects of the addition the KK modes on d_e are weak.

In Fig. 4, we present d_μ with respect to the coupling g for $1/R = 1000 \text{ GeV}$, $\bar{\xi}_{N,\tau\mu}^E = 10 \text{ GeV}$. Here the lower-upper solid (dashed) line represents the EDM, due to the zero mode Higgs-KK mode Higgs included contribution for the parameter $\rho = 0.001$ ($\rho = 0.0005$). Similar to the electron EDM, the sensitivity of d_μ to the coupling g is strong and the addition of KK modes of new Higgs scalars ensures negligible enhancement in the numerical value of d_μ . The muon EDM is at the order of $10^{-23} - 10^{-24} (e - cm)$, for $g \sim 10^{-14}$ and $\rho = 0.001$ ($\rho = 0.0005$). The $1/R$ dependence of d_μ is presented Fig. 5, for $g = 10^{-13}$ and $\bar{\xi}_{N,\tau e}^E = 0.001 \text{ GeV}$. Here the lower-upper solid (dashed) line represents the d_μ , due to the zero mode Higgs-KK mode Higgs included contribution for the parameter $\rho = 0.001$ ($\rho = 0.0005$). The sensitivity of d_μ to $1/R$ is strong similar to the d_e case. If the extra dimension effects are switched off, the muon EDM

⁸Here we choose the masses coming from the Higgs mass term m_S in the lagrangian (see eq. (22)) as $m_{h^0} \sim 0 (GeV)$ and $m_{A^0} \sim 180 (GeV)$ and choose the coupling g in the range such that the zero mode masses $m_{0,h^0(A^0)}$ are around the numerical values as 100 (200) GeV .

d_μ is at the order of $10^{-22} (e - cm)$ and this numerical value can be obtained with the weak localization of the new Higgs doublet, in the present scenario.

Finally, we study the tau EDM d_τ in Fig. 6 and Fig. 7. Fig. 6, represents d_τ with respect to the coupling g for $1/R = 1000 \text{ GeV}$, $\bar{\xi}_{N,\tau\mu}^E = 10 \text{ GeV}$ and $\bar{\xi}_{N,\tau\tau}^E = 50 \text{ GeV}$. Here the lower-upper solid line represents the EDM, due to the zero mode Higgs-KK mode Higgs included contribution for the parameter $\rho = 0.001$. We observe that d_τ enhances up to the values at the order of $2 \times 10^{-22} (e - cm)$ for $g \sim 10^{-14}$ and $\rho = 0.001$. The enhancement due to the KK modes of new Higgs scalars is negligible. Fig. 7 is devoted to the $1/R$ dependence of d_τ for $g = 10^{-13}$ and $\bar{\xi}_{N,\tau e}^E = 0.001 \text{ GeV}$. Here the solid (dashed) line represents the d_τ , due to the zero mode Higgs (KK mode Higgs included) contribution for the parameter $\rho = 0.001$. The sensitivity of d_τ to $1/R$ is strong and almost one order enhancement with the decreasing values of $1/R$ from 1000 (GeV) to 500 (GeV). The tau EDM d_τ is at the order of $10^{-21} (e - cm)$ in the case that the extra dimension effects are switched off. Such a numerical value is obtained for the weak localization of the new Higgs doublet, similar to the electron and muon EDM case.

Now we would like to summarize our results:

- The lepton EDMs are weakly sensitive the KK mode contributions in the given range of the coupling g , however this sensitivity increases for the small values of this coupling. Furthermore, the numerical values of EDMs increase with the decreasing values of the coupling g and increasing values of parameter ρ , which measures the Gaussian widths of leptons in the extra dimension.
- The sensitivity of lepton EDMs to $1/R$ increases with the decreasing values of the parameter ρ .
- We obtain the numerical values at the order of $\sim 10^{-31} (10^{-24}, 10^{-21}) (e - cm)$ for $g \sim 10^{-14}$ and $\rho = 0.001$, for the EDMs $d_e (d_\mu, d_\tau)$, respectively. To be able to reach the results near the current experimental limits, the following domains for the numerical values of the various free parameters should be considered: $\bar{\xi}_{N,\tau e}^E > 0.001 \text{ GeV}$, $\bar{\xi}_{N,\tau\mu}^E > 10 \text{ GeV}$, $\bar{\xi}_{N,\tau\tau}^E > 50 \text{ GeV}$, $1/R < 500 \text{ GeV}$, $\rho > 0.001$ and $g \sim 10^{-14}$. It is obvious that these theoretical values of the EDMs of charged leptons are still far from the experimental limits. Furthermore, the bound for the compactification scale almost coincides with the following bounds existing in the literature: $1/R \sim 300 \text{ GeV}$ [27] in the case of universal extra dimension scenario and $1/R > 500 \text{ GeV}$ from $B \rightarrow \psi K_S$ [19]. Hopefully, the forthcoming

more sensitive experimental measurements will ensure smaller numerical values for the lepton EDMs and, therefore, the possible range of the free parameter set will be more accurate.

As a final word, the future experimental measurements on leptons EDMs would ensure valuable information about the possibility of the existence of localizations of leptons and Higgs bosons in the extra dimensions and make it more clear how the nature behaves beyond the SM.

4 Appendix

4.1 The KK modes of Higgs fields and their masses

Here, we calculate the KK modes of Higgs fields and their masses in the case that the localizer field ϕ_L couples to the new Higgs fields weakly and localize these new fields around $y = 0$, where y is the coordinate of the extra dimension. Our the starting point is the Lagrangian of the new scalar field, which is coupled to the localizer field:

$$L = \sum_{S=h^0, A^0} \left(\frac{1}{2} \partial_M S(x, y) \partial^M S(x, y) - V(S, \phi_L) \right), \quad (21)$$

with $M = 0, 1, \dots, 4$, and

$$\partial_M S(x, y) \partial^M S(x, y) = \partial_\mu S(x, y) \partial^\mu S(x, y) - \left(\frac{dS(x, y)}{dy} \right)^2,$$

with $\mu = 0, 1, 2, 3$. Here we consider the potential $V(S, \phi_L)$,

$$V(S, \phi_L) = \frac{1}{2} m_S^2 S^2(x, y) + \frac{1}{2} g S^2(x, y) \phi_L^2, \quad (22)$$

where ϕ_L is the localizer singlet field, which is chosen as $\phi_L = 2 \mu^2 y$ (see [12]). It has a linear dependence to the new coordinate y for the points near to the fixed point and it couples weakly to new scalar fields $S(x, y)$ ⁹. Now the lagrangian L can be rewritten as

$$L = \frac{1}{2} \partial_\mu S(x, y) \partial^\mu S(x, y) - \frac{1}{2} S(x, y) \left(- \frac{d}{dy^2} + \frac{\partial^2 V}{\partial S^2} \right) S(x, y). \quad (23)$$

With the expansion over the KK modes as

$$S(x, y) = \sum_r h_r(y) S^{(r)}(x), \quad (24)$$

⁹We consider an additional interaction term $\frac{1}{2} g \phi_2^\dagger \phi_2 \phi_L^2$ in the full lagrangian.

one gets the mass equation

$$\left(-\frac{d^2}{dy^2} + m_S^2 + g(2\mu^2)^2 y^2 \right) h_r(y) = m_r^2 h_r(y). \quad (25)$$

The ground state (the zero mode) solution reads

$$h_0(y) = N e^{-\beta y^2}, \quad (26)$$

with the normalization constant $N = \frac{(2\beta)^{1/4}}{\sqrt{\pi^{1/2} \text{Erf}[\pi R \sqrt{2\beta}]}}$ and its corresponding mass $m_{0,S}$

$$m_{0,S}^2 = m_S^2 + 2\beta, \quad (27)$$

with $\beta = \sqrt{g} \mu^2$. Here μ is the energy scale which adjusts the localization of leptons in the extra dimension, namely, $\mu = \frac{1}{\sqrt{2}\sigma}$.

The KK modes for $n \neq 0$ can be obtained by using the operator $D = \frac{d}{dy'} - y'$ as

$$h_r(y) = N^{(r)} D^r e^{-\frac{1}{2} y'^2}, \quad (28)$$

where $y' = \sqrt{2\beta} y$ and $N^{(r)}$ is the normalization constant for r_{th} KK mode,

$N^{(r)} = \frac{\beta^{1/4}}{\sqrt{\pi^{1/2} 2^{n-\frac{1}{2}} (n)! \sqrt{\text{Erf}[\pi R \sqrt{2\beta}]}}}$. Finally the masses of the KK modes, including the zero one, becomes ¹⁰

$$m_{r,S}^2 = m_S^2 + 2\beta(2r+1). \quad (29)$$

¹⁰Notice that we take even KK modes of new Higgs bosons, namely the ones with $r = 2n$, $n = 0, 1, \dots$, since the extra dimension is compactification on the orbifold S^1/Z_2 . In this case, the KK mode masses read $m_{r,S}^2 = m_S^2 + 2\beta(4n+1)$

References

- [1] C. R. Schmidt and M. E. Peskin, *Phys. Rev. Lett.* **69** (1992) 410.
- [2] E. D. Commins et.al. *Phys. Rev. A* **50** (1994) 2960.
- [3] J. Bailey et al, *Journ. Phys.* **G4** (1978) 345;
- [4] Particle Data Group, D. E. Groom et.al. *European Phys. Journ.* **C15** (2000) 1.
- [5] B. Dutta, R. N. Mohapatra, *Phys. Rev.* **D68** (2003) 113008.
- [6] E. Iltan, *Phys. Rev.* **D64** (2001) 013013.
- [7] E. Iltan, *JHEP* **065** (2003) 0305.
- [8] E. Iltan, *JHEP* **0404** (2004) 018.
- [9] E. Iltan, *Eur. Phys.* **C44** (2005) 411.
- [10] E. P. Sahabalin, *Sov. J. Nucl. Phys.* **28** (1978) 75; J. F. Donoghue, *Phy. Rev.* **D18** (1978) 1632; A. Czarnecki and B. Krause, *Phys. Rev. Lett.* **78** (1997) 4339; I. B. Khriplovich, *Yad. Fiz.* **44** (1986) 1019 (*Sov. J. Nucl. Phys.* **44** (1986) 659); *Phys. Lett.* **B173** (1986) 193; A. Czarnecki and B. Krause, *Acta Phys. Polon.* **B28** (1997) 829; *Phys. Rev. Lett* **78** (1997) 4339; S. Weinberg, *Phys. Rev. Lett.* **58** (1976) 657; N. G. Deshpande and E. Ma, *Phys. Rev.* **D 16** (1977) 1583; for a recent review, see H. Y. Cheng, *Int. J. Mod. Phys.* **A 7** (1992) 1059; G. C. Branco and M. N. Rebelo, *Phys. Lett.* **B 160** (1985) 117; J. Liu and L. Wolfenstein, *Nucl. Phys.* **B 289** (1987) 1; C. H. Albright, J. Smith and S. H. H. Tye, *Phys. Rev. D.* **D 21** (1980) 711; D. Atwood, S. Bar-Shalom, A. Soni, *Phys. Rev. D* **51** (1995) 1034; E. Iltan, *J. Phys.* **G27** (2001) 1723; E. Iltan, *Phys. Rev.* **D65** (2002) 073013; Y. Liao and X. Li, *Phys. Rev.* **D60** (1999) 073004; D. G. Dumm and G. A. G Sprinberg,, *Eur-Phys. J.* **C 11** (1999) 293; A. Soni and R. M. Xu, *Phys. Rev. Lett.* **69** (1992) 33.
- [11] V. Baryshevsky, hep-ph/0402209; O. Lebedev, K. A. Olive, M. Pospelov, A. Ritz, *Phys. Rev.* **D70** (2004) 016003; A. G. Rodriguez, M. A. H. Ruiz, A. Santiago, *Phys. Rev.* **D69** (2004) 073008; A. Gutierrez-Rodriguez, M.A.Hernandez-Ruiz and L.N. Luis-Noriega, *Mod. Phys. Lett.* **A19** (2004) 2227; R. E. Mischke, AIP Conf.Proc.675:246-250,2003; D. A. Demir, Oleg Lebedev, K. A. Olive, M. Pospelov, A. Ritz, *Nucl.Phys.* **B680**, (2004) 339; M. Endo, M. Kakizaki, M. Yamaguchi, *Phys.Lett.* **B 583** (2004) 186; D. Kawall, F. Bay, S. Bickman,

- Y. Jiang, D. DeMille, *Phys. Rev. Lett.* **92** (2004) 133007; I. Masina, C. Savoy, *Phys.Lett.* **B579** (2004) 99; J. Hisano, Y. Shimizu, *Phys.Lett.* **B581** (2004) 224-230; A. Syntfeld, H. Mach, W. Kurcewicz, B. Fogelberg, W. Pociennik, E. Ruchowska *Phys.Rev.* **C68** (2003) 024304; T. Suzuki, H. Sagawa, K. Hagino, *Phys.Rev.* **C68** (2003) 014317.
- [12] N. Arkani-Hamed and M. Schmaltz, *Phys. Rev.* **D61**, 033005 (2000); N. Arkani-Hamed, Y. Grossman and M. Schmaltz, *Phys. Rev.* **D61** (2000) 115004.
- [13] I. Antoniadis, *Phys. Lett.* **B246**, 377 (1990).
- [14] E. A. Mirabelli, Schmaltz, *Phys. Rev.* **D61** (2000) 113011.
- [15] W. F. Chang, I. L. Ho and J. N. Ng, *Phys. Rev.* **D66** 076004 (2002).
- [16] G. C. Branco, A. Gouvea, M. N. Rebelo, *Phys. Lett.* **B506** (2001) 115.
- [17] W. F. Cang, J. N. Ng, *JHEP* **0212** (2002) 077.
- [18] Y. Grossman and G. Perez, *Phys. Rev.* **D67** (2003) 015011 ; *Pramana* **62** (2004) 733.
- [19] B. Lillie, J. L. Hewett, *Phys. Rev.* **68** (2003) 116002.
- [20] A. Delgado, A. Pomarol, M. Quiros, *JHEP* **0001** (2000) 030; Y. Grossman, *Int. J. Mod. Phys.* **A15** (2000) 2419; D. E. Kaplan and T. M. Tait, *JHEP* **0111** (2001) 051; G. Barenboim, *et. al.*, *Phys. Rev.* **D64** (2001) 073005; W. F. Chang, I. L. Ho and J. N. Ng, *Phys. Rev.* **D66** (2002) 076004; W. Skiba and D. Smith, *Phys. Rev.* **D65** (2002) 095002; Y. Grossman, R. Harnik, G. Perez, M. D. Schwartz and Z. Surujon, *Phys. Rev.* **D71** (2004) 056007; P. Dey, G. Bhattacharya, *Phys. Rev.* **D70** (2004) 116012; Y. Nagatani, G. Perez, *JHEP* **0502**(2005) 068; R. Harnik, G. Perez, M. D. Schwartz, Y. Shirman, **0503** (2005) 068.
- [21] E. O. Iltan, hep-ph/0504013.
- [22] E. O. Iltan, *Eur.Phys.J.* **C46** (2006) 487.
- [23] E. O. Iltan, *Eur.Phys.J.* **C46** (2006) 475.
- [24] Z. Surujon, *Phys. Rev.* **D73** (2006) 016008.
- [25] M. Deile, et. al., Muon g-2 Collaboration, , *Nucl.Phys.Proc.Suppl.* **116** (2003) 215.
- [26] E. Iltan, H. Sundu, *Acta Phys.Slov.* **53** (2003) 17.

- [27] T. Appelquist, H.-C. Cheng, B. A. Dobrescu, *Phys. Rev.* **D64** 035002 (2001); T. Appelquist, H. Yee, *Phys. Rev.* **D67** 055002, 2003;
- [28] T. G. Rizzo, J. D. Wells, *Phys. Rev.* **D61**, 016007 (2000).




Figure 1: One loop diagrams contribute to EDM of charged leptons due to neutral Higgs bosons h^0 , A^0 in the 2HDM, including KK modes in a single extra dimension. Wavy lines represent the electromagnetic field and dashed lines the Higgs field where $l_{1(i)} = e, \mu, \tau$




Figure 2: d_e with respect to the coupling g for $1/R = 1000 \text{ GeV}$, $\bar{\xi}_{N,\tau e}^E = 0.001 \text{ GeV}$. Here the lower-upper solid (dashed) line represents the EDM, due to the zero mode Higgs-KK mode Higgs included contribution for the parameter $\rho = 0.001$ ($\rho = 0.0005$).




Figure 3: d_e with respect to $1/R$, for $g = 10^{-13}$ and $\bar{\xi}_{N,\tau e}^E = 0.001 \text{ GeV}$. Here the lower-upper solid (dashed) line represents the d_e , due to the zero mode Higgs-KK mode Higgs included contribution for the parameter $\rho = 0.001$ ($\rho = 0.0005$).




Figure 4: The same as Fig. 2 but for d_μ and $\bar{\xi}_{N,\tau\mu}^E = 10 \text{ GeV}$.




Figure 5: The same as Fig. 3 but for d_μ and $\bar{\xi}_{N,\tau\mu}^E = 10 \text{ GeV}$.




Figure 6: The same as Fig. 2 but for d_τ and $\rho = 0.001$, $\bar{\xi}_{N,\tau\mu}^E = 10 \text{ GeV}$ and $\bar{\xi}_{N,\tau\tau}^E = 50 \text{ GeV}$.




Figure 7: The same as Fig. 3 but for d_τ and $\rho = 0.001$, $\bar{\xi}_{N,\tau\mu}^E = 10 GeV$ and $\bar{\xi}_{N,\tau\tau}^E = 50 GeV$.